

## University of Groningen

### Aortasclerose : Bevindingen bij een bevolkingsonderzoek te Zutphen

Schermers, Govert Gerrit

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

1969

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Schermers, G. G. (1969). *Aortasclerose : Bevindingen bij een bevolkingsonderzoek te Zutphen*. [, Rijksuniversiteit Groningen]. [S.n.].

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## STELLINGEN

### I

Aortasclerose is een, op oudere leeftijd, regelmatig voorkomende aandoening, welke van aortastenose onderscheiden dient te worden.

### II

Bij patienten met een verwijde aortaboog en kalkafzettingen in de wand hiervan moet de diagnose aortasclerose overwogen worden.

### III

De revalidatie van patienten met cardiovasculaire afwijkingen kan efficiënter worden door inschakeling van een ergonoom.

### IV

Elke patient met lumbago die veertien dagen na het begin van zijn klachten geen duidelijke verbetering vertoont dient tenminste eenmaal specialistisch te worden onderzocht.

### V

Bij optimale verloskunde van de huisarts behoort het bijhouden van baringsverslagen en een voorlichtingsgesprek met de primigravidae en haar echtgenoten, enige weken voor de berekende datum van de partus.

### VI

Het gebruik van de bekkenpasser voor verloskunde door de huisarts is slechts van academisch belang.

### VII

Bij het onderzoek van de lumbosacrale wervelkolom of het bekken naar skeletmetastasen met hulp van Strontium-85 dient rekening gehouden te worden met het aanwezig zijn van het isotoop in de faeces.

Het opsporen van tumormetastasen in het skelet met hulp van Strontium-85.

D. J. Dronkers, acad. proefschrift 1966.

---

## VIII

Bij radiotherapie is de fractiedosis in eerste instantie bepalend voor de uiteindelijke reactie van zowel het normale als het tumorweefsel.

Dr. G. Kok, voordracht 192e alg. ledenverg. Ned. Ver.  
voor Radiologie, Dec. 1968.

## IX

Het staat niet vast, dat door ijzerdepletie (sideropenie, latent ijzergebrek) klachten veroorzaakt worden.

## X

In de *stercorarius*groep van het genus *coprinus* behoren de ronde velumcellen niet spherocysten, maar spherocyten genoemd te worden.

The 'stercorarius group' of the genus *coprinus*, E. Kits  
van Waveren, *Persoonia* 1968, Vol. 5, Part 2, pp.  
131—176.